



1



4



2

malaria how big a problem...

- Most malaria cases in 2017 were in the WHO African Region (200 million or 92%), followed by the WHO South-East Asia Region with 5% of the cases and the WHO Eastern Mediterranean Region with 2%.
- Fifteen countries in sub-Saharan Africa and India carried almost 80% of the global malaria burden. Five countries accounted for nearly half of all malaria cases worldwide: Nigeria (25%), Democratic Republic of the Congo (11%), Mozambique (5%), India (4%) and Uganda (4%).
- Nearly 80% of global malaria deaths in 2017 were concentrated in 17 countries in the WHO African Region and India; 7 of these countries accounted for 53% of all global malaria deaths: Nigeria (19%).

Chevron
human energy

5



3

WORKFORCE DEMOGRAPHICS


- Nationals
- Expatriates
 - Dependents
- Rotators

Chevron
human energy

6

Chevron Global Health and Medical

- Americas
 - Occupational Health
 - Occupational health physicians
 - Nurse Practitioners, Nurses
 - EAP
 - Public Health
- EEAME (Europe, Eurasia, Africa, Middle East)
 - Hospitals
 - Clinics
 - Able to see Nationals and Expatriates
 - Onsite Hospitals
- Asia Pacific
 - Hospitals
 - Clinics
 - Able to see Nationals and Expatriates
 - Onsite Hospitals





7


Chevron Malaria Hotline




Malaria Hotline

- Fever
- Headache
- Sweats and chills
- Pain in the back and joints
- Fatigue
- Cough
- Diarrhea
- Nausea
- Vomiting

If you or your medical provider have questions, call the Chevron Malaria Hotline at **+1-866-276-5118**.

If you have visited a malarious area within the last six months and are experiencing any of these symptoms, go to a hospital or doctor immediately to be tested for malaria.






10

Chevron Guidelines for travelers

- Visit or call a Chevron medical clinic two weeks before traveling to a malaria-endemic area
- Complete mosquito-borne illness training before departure and every 2 years
- Begin taking anti-malaria medications before the trip
- Anti-malaria drugs that currently are in use are: Malarone®, Doxycycline and Lariam.
- Continue taking anti-malaria medications after leave the area

8

Malaria testing kit








9